

A GRASSMANN REPRESENTATION OF THE HUBBLE PARAMETER

HARET C. ROSU and JOSE TORRES

Instituto de Física, Universidad de Guanajuato, Apdo Postal E-143, 37150 León, Gto, Mexico

The Riccati equation for the Hubble parameter H of barotropic FRW cosmologies in conformal time for $\kappa \neq 0$ spatial geometries and in comoving time for the $\kappa = 0$ geometry, respectively, is generalized to odd Grassmannian time parameters. We obtain a system of simple differential equations for the four supercomponents (two of even type and two of odd type) of the Hubble superfield function $\mathcal{H}$ that is explicitly solved. The second even Hubble component does not have an evolution governed by general relativity although there are effects of the latter upon it.

1 - Recently, Faraoni,¹ showed that the equations describing barotropic FRW cosmologies can be combined in a simple Riccati equation leading in a much easier way to the textbook solutions for the FRW scale factors. Faraoni obtained the following cosmological Riccati equation

$$\frac{dH}{d\eta} = -cH^2 - \kappa c, \quad (1a)$$

for the log derivative of the FRW scale factor, the famous Hubble parameter $H(\eta) = \frac{da/d\eta}{a}$. The independent variable is the conformal time η , $c = \frac{3}{2}\gamma - 1$ was assumed constant by Faraoni, and $\kappa = 0, \pm 1$ is the curvature index of the flat, closed, open FRW universe, respectively. The adiabatic index γ is defined through the barotropic equation of state $P = (\gamma - 1)\rho$, where P and ρ are, respectively, the pressure and energy density of the cosmological fluid under consideration. It is a constant quantity in the approach of Faraoni. It is a simple matter to write the first Friedmann dynamical equation $\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\rho + P)$ in the form $\frac{\ddot{a}}{a} = -\frac{4\pi G\rho}{3}c$ for the barotropic case. Thus, the common lore is that a negative c ($\gamma < \frac{2}{3}$) implies an accelerating universe as claimed in the astrophysics of supernovae.

We also notice that from the mathematical point of view, Faraoni's FRW Riccati equation being of constant coefficients is directly integrable. The solutions are

$$H^+ = -\tan c\eta, \quad H^- = \coth(c\eta), \quad (1b)$$

for the closed and open FRW universes, respectively.

In the following we perform a very simple generalization of the cosmological Riccati equation (1a) by considering the Hubble parameter as a function not only of time but also of the so-called odd "time" parameters, θ_1 and $\bar{\theta}_1$ (where $\bar{\theta}_1$ is the complex conjugate of θ_1).² In this way, the Hubble parameter becomes a superfield

in the terminology of superanalysis.³ This is only a toy generalization since we do not relate it to any sophisticated formalism, such as supergravity. However, we obtain some simple analytic formulas for all the components of the Hubble superfield. We mention that similar types of supersymmetric extensions (affecting in addition the operatorial structure) are of much interest in the theory of integrable models that appear naturally in the study of strings in the matrix model approach.⁴

2 - We shall write the Riccati equation for the superfield $\mathcal{H}$ in the following form

$$\frac{d\mathcal{H}}{d\eta} \equiv \frac{i}{2} \{D_{\theta_1}, D_{\bar{\theta}_1}\} \mathcal{H} = \alpha \mathcal{H}^2 + \beta, \quad (2)$$

where

$$\mathcal{H}(\eta, \theta_1, \bar{\theta}_1) = H_0(\eta) + i\theta_1 \bar{H}_1(\eta) + i\bar{\theta}_1 H_1(\eta) + \theta_1 \bar{\theta}_1 H_{11}(\eta), \quad (3)$$

$\alpha = -c$, $\beta = -\kappa c$, and

$$D_{\theta_1} = \frac{d}{d\theta_1} + i\bar{\theta}_1 \frac{d}{d\eta}, \quad D_{\bar{\theta}_1} = -\frac{d}{d\bar{\theta}_1} - i\theta_1 \frac{d}{d\eta}. \quad (4)$$

The superfield $\mathcal{H}$ satisfies the reality condition $\mathcal{H}^\dagger = \mathcal{H}$ and is assumed to be dimensionless implying the following η dimensions

$$[\eta] = 1, \quad [\theta_1] = 1/2, \quad [H_0] = 0, \quad [H_1] = -1/2, \quad [H_{11}] = -1.$$

After simple (super)calculations we get the following system of equations

$$\frac{dH_0}{d\eta} = \alpha H_0^2 + \beta, \quad (5a)$$

$$\frac{d\bar{h}_1}{d\eta} = 2\alpha H_0 \bar{h}_1, \quad (5b)$$

$$\frac{dh_1}{d\eta} = 2\alpha H_0 h_1, \quad (5c)$$

$$\frac{dH_{11}}{d\eta} = 2\alpha(H_0 H_{11} + \bar{h}_1 h_1 \bar{\epsilon} \epsilon), \quad (5d)$$

where $\bar{h}_1 \bar{\epsilon} = \bar{H}_1$, $h_1 \epsilon = H_1$, and ϵ is a Grassmann parameter.

One can see that equation (5a) is identical to Faraoni's Riccati equation. Therefore, the solutions are

$$H_0^+ = -\tan(c\eta), \quad H_0^- = \coth(c\eta), \quad (6)$$

For the other components, we get simple first order differential equations. The equations (5b) and (5c) are identical since we consider only the possibility of real superpartners. The solutions are easily obtained

$$h_1^+ = \frac{1}{\cos^2(c\eta)}, \quad h_1^- = \frac{1}{\sinh^2(c\eta)}, \quad (7)$$

for the closed and open spatial geometries, respectively.

Finally, the solution of the equation for H_{11} can be obtained by the method of the variation of constants leading to

$$H_{11}^{\pm} = e^{2\alpha \int_{\eta_0}^{\eta} H_0 d\eta} \left[A_{11}^{\pm} + \bar{\epsilon}\epsilon \int_{\eta_0}^{\eta} 2h_1^2 e^{-2\alpha \int_{\eta_0}^{\eta} H_0 d\eta} d\eta \right], \quad (8)$$

where the parameters $A_{11}^{\pm}$ are related to the initial conditions. We thus obtain

$$H_{11}^{+} = \cos^{-2}(c\eta) (A_{11}^{+} - 2\bar{\epsilon}\epsilon \tan(c\eta)) \quad (9)$$

and

$$H_{11}^{-} = \sinh^{-2}(c\eta) (A_{11}^{-} + 2\bar{\epsilon}\epsilon \coth(c\eta)) . \quad (10)$$

3 - The case $\kappa = 0$ is special in the sense that $\alpha = -(c+1)$, $\beta = 0$ and the independent variable is the comoving time t .⁵ Therefore, the odd time parameters refer to the comoving time in this case.

(i) For $c+1 \neq 0$ ($\alpha \neq 0$, $\beta = 0$) we get

$$H_0^0 = \frac{1}{c+1} \frac{1}{t}, \quad h_1^0 = \bar{h}_1^0 = \frac{1}{t^2}, \quad H_{11}^0 = A_{11}^0 t^{-2} + 2\bar{\epsilon}\epsilon(c+1)t^{-3} . \quad (11)$$

(ii) The case $c+1 = 0$ ($\alpha = 0$, $\beta = 0$) leads to

$$H_0 = C_0, \quad h_1 = \bar{h}_1 = C_1, \quad H_{11} = C_{11} . \quad (12)$$

4 - Through a simple generalization of the barotropic FRW Riccati evolution equation to odd Grassmann time parameters $(\theta_1, \bar{\theta}_1)$ a Grassmann (superfield) representation of the Hubble parameter is given in this work. The superfield components are explicitly obtained under the simple assumption of a real Hubble superfield. We notice that the evolution equation of the second even component H_{11} is a linear first order differential equation not a nonlinear Riccati equation as required by Friedmann equations. Thus, we are out of the realm of general relativity although there is an induced effect of the latter as seen from the presence of H_0 in equation (5d).

Acknowledgements

The authors thank Prof. V. Tkach and Dr. J.J. Rosales for discussions.

References

1. V. Faraoni, Am. J. Phys. **67**, 732 (1999) [physics/9901006]. See also, J.A.S. Lima, Am. J. Phys., to appear, [astro-ph/0109215]. For nonrelativistic supersymmetric approaches with application to cosmological acceleration, see H.C. Rosu, Mod. Phys. Lett. A **15**, 979 (2000) [gr-qc/0003108].
2. V.I. Tkach, J.J. Rosales and J. Socorro, Mod. Phys. Lett. A **14**, 1209 (1999) and the references therein.
3. F.A. Berezin, *Introduction to Superanalysis*, Reidel, Dordrecht, 1987.
4. A. Das, *Integrable Models*, World Scientific, 1989; For recent work, see e.g., A. Das and Z. Popowicz, Phys. Lett. A **274**, 30 (2000) and references therein; R. Jackiw, hep-th/0110057 and physics/0010079; see also, A. Turbiner, hep-th/9209080.
5. H.C. Rosu, Mod. Phys. Lett. A **16**, 1147 (2001) [gr-qc/0003108].